

Cancer Association of South Africa (CANSA)



Fact Sheet on the Amount of Ionizing Radiation Received from Routine Imaging

Introduction

Ionizing radiation is any form of radiation with enough energy to break off electrons from atoms (that is, to ionize the atoms). This radiation can break the chemical bonds in molecules, including DNA molecules, thereby disturbing their normal functioning. X-rays and gamma rays are the only common forms of radiation with sufficient energy to penetrate and damage body tissue below the surface of the skin.

[Picture Credit: Ionizing Radiation]



Among the many sources of ionizing radiation are traditional X-rays, computed tomography (CT) scans, fluoroscopy, and other medical radiological procedures. A newer source of X-rays is the use of backscatter scanners in airport security. Sources of gamma rays include emissions from nuclear power plants, scientific research involving radionuclides, military weapons testing, and nuclear medicine procedures such as bone, thyroid and lung scans (Brenner, 2011).

Key Facts on Ionizing Radiation

According to the World Health Organization (WHO):

- Ionizing radiation is a type of energy released by atoms in the form of electromagnetic waves or particles
- People are exposed to natural sources of ionizing radiation, such as in soil, water, and vegetation, as well as in human-made sources, such as X-rays and medical devices
- Ionizing radiation has many beneficial applications, including uses in medicine, industry, agriculture and research
- As the use of ionizing radiation increases, so does the potential for health hazards if not properly used or contained
- Acute health effects such as skin burns or acute radiation syndrome can occur when doses of radiation exceed certain levels

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]


October 2018

Page 1

- Low doses of ionizing radiation can increase the risk of longer term effects such as cancer

Types of Ionizing Radiation

Ionizing radiation takes a few forms: Alpha, beta, and neutron particles, and gamma and X-rays. All types are caused by unstable atoms, which have either an excess of energy or mass (or both). In order to reach a stable state, they must release that extra energy or mass in the form of radiation.



(Mirion Technologies).

Radiation-Related Quantities

The following table shows radiation quantities in SI and non-SI units

Quantity	Name	Symbol	Unit
Exposure (X)	roentgen	R	esu / 0.001293 g of air
Absorbed dose (D)	Rad	rad	erg·g ⁻¹
	gray	Gy	J·kg ⁻¹
Activity (A)	curie	Ci	3.7 × 10 ¹⁰ s ⁻¹
	becquerel	Bq	s ⁻¹
Dose equivalent (H)	Roentgen equivalent man	rem	100 erg·g ⁻¹
	sievert	Sv	J·kg ⁻¹
Fluence (Φ)	(reciprocal area)		cm ⁻² or m ⁻²

(Wikipedia)

Risks of Ionizing Radiation

The unit of measurement used for the biological effect of radiation on the human body is the millisievert (mSv). The average global exposure to natural radiation is 2.4 mSv per year. It is known that very large doses of over 5 000 mSv, received by the entire body over a short time, result in death within a few days. We know, however, that some of the effects of exposure to radiation do not appear unless a certain large dose has been absorbed. Doses over 100 mSv can have a harmful effect on humans, such as a higher incidence of developing cancer.

At even lower doses of radiation, below 100 mSv, there is a lot of uncertainty about the overall effects. What we do know is that the risk of adverse effects in this dose range is very low. To be on the safe side, we assume that there is a risk even in this low dose range and this risk is proportional to the dose by the same amount as in the high dose ranges.

Statistically, everyone has a one in three chance of developing cancer at some point in their life. In order to put the risk of cell damage caused by radiation exposure in medical imaging into some perspective, the UK Health Protection Agency (HPA) has calculated that:

- an X-ray of one's chest, teeth, arms or feet is the equivalent to a few days' worth of background radiation and has a less than one in a million chance of causing cancer,
- an X-ray of one's skull or neck is the equivalent to a few weeks' worth of background radiation and has a 1 in 100 000 to 1 in 1 000 000 chance of causing cancer,
- an X-ray of one's breasts (mammogram), hip, spine, abdomen or pelvis is the equivalent of a few months' to a year's worth of background radiation and has a 1 in 10 000 to 1 in 100 000 chance of causing cancer, and
- an X-ray that uses a contrast fluid, such as a barium meal, is the equivalent of a few years' worth of background radiation and has a 1 in 1 000 to 1 in 10 000 chance of causing cancer.


(Understanding Medical Radiation).

Radiation and DNA

Radiation is simply a mechanism whereby energy passes through space. It takes the form of an electromagnetic wave, with the frequency of the electromagnetic wave determining its position in the electromagnetic spectrum. Low-frequency waves such as radio waves lie at one end of the spectrum and high-energy, high-frequency X-rays/Gamma rays at the other end.

[Picture Credit: Electromagnetic Spectrum]

These high-frequency, high-energy waves, are termed "ionizing" (as opposed to non-ionizing) radiation because they contain sufficient energy to displace an electron from its orbit around a nucleus. The most important consequence of this displaced electron on human tissue is the potential damage it can inflict on DNA, which may occur directly or indirectly. Direct damage occurs when the



Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

October 2018

displaced electron hits and breaks a DNA strand. Indirect damage occurs when the electron reacts with a water molecule, creating a powerful hydroxyl radical which then damages the cell's DNA.

Damage to a cell's DNA in either of these ways can have several consequences. A single-strand DNA break is usually repaired appropriately by the cell with no subsequent deleterious sequelae. However, a break affecting both strands of DNA allows the potential for abnormal reconnection of the strands, which likely accounts for all the adverse biological effects ionizing radiation has on humans.

First, DNA may rejoin itself incorrectly, rendering the cell nonviable with cell death (apoptosis) taking place. Second, it may rejoin as a symmetrical translocation with the potential expression of an oncogene during division (and development of subsequent malignancy) or with abnormal division in gonads, giving rise to potential hereditary disorders.

Radiosensitivity is the probability of a cell, tissue, or organ suffering an effect per unit dose of radiation. Radiosensitivity is highest in cells which are highly mitotic or undifferentiated. For this reason the basal epidermis, bone marrow, thymus, gonads, and lens cells are highly radiosensitive. Muscle, bones, and nervous system tissues have a relative low radiosensitivity. (Goodman).

Health Effects of Ionizing Radiation

Radiation damage to tissue and/or organs depends on the dose of radiation received, or the absorbed dose which is expressed in a unit called the gray (Gy). The potential damage from an absorbed dose depends on the type of radiation and the sensitivity of different tissues and organs.

The *effective dose* is used to measure ionizing radiation in terms of the potential for causing harm. The sievert (Sv) is the unit of effective dose that takes into account the type of radiation and sensitivity of tissues and organs. It is a way to measure ionizing radiation in terms of the potential for causing harm. The Sv takes into account the type of radiation and sensitivity of tissues and organs.

The Sv is a very large unit so it is more practical to use smaller units such as millisieverts (mSv) or microsieverts (μ Sv). There are one thousand μ Sv in one mSv, and one thousand mSv in one Sv. In addition to the amount of radiation (dose), it is often useful to express the rate at which this dose is delivered (dose rate), such as microsieverts per hour (μ Sv/hour) or millisievert per year (mSv/year).

Beyond certain thresholds, radiation can impair the functioning of tissues and/or organs and can produce acute effects such as skin redness, hair loss, radiation burns, or acute radiation syndrome. These effects are more severe at higher doses and higher dose rates. For instance, the dose threshold for acute radiation syndrome is about 1 Sv (1 000 mSv).

If the radiation dose is low and/or it is delivered over a long period of time (low dose rate), the risk is substantially lower because there is a greater likelihood of repairing the damage. There is still a risk of long-term effects such as cancer, however, that may appear years or even decades later. Effects of this type will not always occur, but their likelihood is proportional to the radiation dose. This risk is higher for children and adolescents, as they are significantly more sensitive to radiation exposure than adults.

Epidemiological studies on populations exposed to radiation, such as atomic bomb survivors or radiotherapy patients, showed a significant increase of cancer risk at doses above 100 mSv. More recently, some epidemiological studies in individuals exposed to medical exposures during childhood (paediatric CT) suggested that cancer risk may increase even at lower doses (between 50-100 mSv).

Prenatal exposure to ionizing radiation may induce brain damage in foetuses following an acute dose exceeding 100 mSv between weeks 8-15 of pregnancy and 200 mSv between weeks 16-25 of pregnancy. Before week 8 or after week 25 of pregnancy human studies have not shown radiation risk to foetal brain development. Epidemiological studies indicate that the cancer risk after foetal exposure to radiation is similar to the risk after exposure in early childhood.

(World Health Organization).

Other Effects of Ionizing Radiation

The deleterious effect ionizing radiation has on human tissue can be divided into two types: non-stochastic (deterministic) or stochastic effects.

Deterministic (Non-Stochastic) Effects - Deterministic effects only occur once a threshold of exposure has been exceeded. The severity of deterministic effects increases as the dose of exposure increases. Because of an identifiable threshold level, appropriate radiation protection mechanisms and occupational exposure dose limits can be put in place to reduce the likelihood of these effects occurring.

Deterministic effects are caused by significant cell damage or death. The physical effects will occur when the cell death burden is large enough to cause obvious functional impairment of a tissue or organ.

Examples

1.

Skin erythema/necrosis/epilation - Erythema occurs 1 to 24 hours after 2 Sv have been received. Breakdown of the skin surface occurs approximately four weeks after 15 Sv have been received. Epilation is reversible after 3 Sv but irreversible after 7 Sv and occurs three weeks following exposure.

2.

Cataract - Cataract occurs due to accumulation of damaged or dead cells within the lens, the removal of which cannot take place naturally. Cataract occurs after 2 to 10 Gy have been received, but may take years to develop.

3.

Sterility - Radiation can impair oocyte function, leading to impaired or non-fertility. The radiation dose required to have this effect decreases with age due to falling total oocyte numbers. Similarly, radiation exposure to the testes can result in temporary or permanent azoospermia. Permanent sterility occurs after 2.5 to 3.5 Gy have been received by the gonads.

4.

Radiation Sickness – Radiation sickness (correctly termed acute radiation syndrome) involves nausea, vomiting, and diarrhoea developing within hours or minutes of a radiation exposure. This is due to deterministic effects on the bone marrow, GI tract, and CNS.

5.

IUGR/Teratogenesis/Foetal Death - Deterministic radiation exposure effects during pregnancy depend not only on the radiation dose received but also on the gestational age at which it occurred. The embryo is relatively radio-resistant during its preimplantation phase but highly radiosensitive during its organogenesis (2 to 8 weeks) and neuronal stem cell proliferation phases (8 to 15 weeks). Foetal radiosensitivity falls after this period. High levels of radiation exposure in pregnancy can lead to growth retardation, in particular microcephaly. The threshold dose for this effect is high (>20Gy) with other deterministic effects (hypospadias, microphthalmia, retinal degeneration, and optic atrophy) having a lower threshold level of >1Gy.

Stochastic Effects - Current thinking is that stochastic effect occurrence follows a linear no-threshold hypothesis. This means that although there is no threshold level for these effects, the risk of an effect occurring increases linearly as the dose increases.

Examples

1.

Cancer - Over time, anecdotal evidence suggested that ionizing radiation could cause cancer. However, reliable evidence has only relatively recently become available. Data from the Radiation Effects Research Foundation on individuals exposed to radiation from the atomic bombs in Hiroshima and Nagasaki have shown an increased relative risk of developing malignancy (leukaemia, oral cavity, oesophagus, stomach, colon, lung, breast, ovary, urinary bladder, thyroid, liver, non-melanoma skin, and nervous system) as a result of radiation exposure. As such, multiple bodies, including the U.S. Department of Health and Human Services, have classified ionizing radiation as a human carcinogen.

Unfortunately, doses which have been shown to result in this increased relative malignancy risk are similar to levels which can also be imparted by radiology studies such as CT scans, interventional radiology, and barium enema procedures. Indeed, the excessive relative risk of cancer mortality determined by the International Commission of Radiological Protection (ICRP) is 5%/Sv. The National Research Council of the National Academies has concluded that a single CT scan with a dose of 10 mSv carries a risk of 1:1 000 of producing cancer. A comprehensive list of effective doses in various other radiology and nuclear medicine procedures can be easily compared with natural background radiation levels.

The risk of developing solid cancers follows a linear pattern with increasing dose although the age at which exposure takes place is highly relevant. A decline in radiosensitivity does take place with age, making young children more susceptible to radiation-induced malignancies. Although the malignancy risk for the population as a whole is 5%/Sv, this rises to 15%/Sv in a young girl and falls to 1%/Sv in a 70-year-old.

Adopting the linear no-threshold hypothesis and extrapolating the data to very low dose exposures (plain radiographs, for example) would suggest that while the risk of a radiation-induced malignancy persists, its impact becomes negligible for a single exposure.

There is some controversy concerning the extrapolation of the linear no-threshold hypothesis to very low doses given that no increased incidence of cancer is seen in areas of high background radiation or in airline pilots. Caution should be exercised in multiple radiation exposures, however, given that stochastic effects are cumulative. Similarly, multiple high-dose diagnostic imaging procedures such as CT can easily exceed the levels known to impart an increased relative risk for malignancy.

2.

Hereditary Defects (e.g., Down Syndrome) - Although the incidence of hereditary defects in patients exposed to radiation in Japan and Chernobyl have shown no increased evidence for hereditary defects, animal experiments would suggest that this risk does exist. The United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) and ICRP propose a hereditary defect risk of between 0.3 to 0.8% per Sv.

(Image Wisely).

Effective Radiation Received from Selected Imaging Procedures

The following is an indication of the effective radiation a person will normally receive from certain selected imaging procedures.

EFFECTIVE RADIATION DOSE IN ADULTS FOR THIS PROCEDURE	Adult approximate effective radiation dose:
Abdominal Region	
Computed Tomography (CT) - Abdomen and Pelvis	10 mSv
Computed Tomography (CT) - Abdomen and Pelvis, repeated with and without contrast material	20 mSv
Computed Tomography (CT) - Colonography	6 mSv
Intravenous Pyelogram (IVP)	3 mSv
Radiography (X-ray) - Lower GI Tract	8 mSv
Radiography (X-ray) - Upper GI Tract	6 mSv

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

October 2018

Bone	
Radiography (X-ray) - Spine	1.5 mSv
Radiography (X-ray) - Extremity	0.001 mSv
Central Nervous System	
Computed Tomography (CT) - Head	2 mSv
Computed Tomography (CT) - Head, repeated with and without contrast material	4 mSv
Computed Tomography (CT) - Spine	6 mSv
Chest	
Computed Tomography (CT) - Chest	7 mSv
Computed Tomography (CT) - Lung Cancer Screening	1.5 mSv
Radiography - Chest	0.1 mSv
Dental	
Intraoral X-ray	0.005 mSv
Heart	
Coronary Computed Tomography Angiography (CTA)	12 mSv
Cardiac CT for Calcium Scoring	3 mSv
Men's Imaging	
Bone Densitometry (DEXA)	0.001 mSv
Nuclear Medicine	
Positron Emission Tomography – Computed Tomography (PET/CT)	25 mSv
Women's Imaging	
Bone Densitometry (DEXA)	0.001 mSv
Mammography	0.4 mSv

(RadiologyInfo).

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

October 2018

Medical Disclaimer

This Fact Sheet is intended to provide general information only and, as such, should not be considered as a substitute for advice, medically or otherwise, covering any specific situation. Users should seek appropriate advice before taking or refraining from taking any action in reliance on any information contained in this Fact Sheet. So far as permissible by law, the Cancer Association of South Africa (CANSA) does not accept any liability to any person (or his/her dependants/estate/heirs) relating to the use of any information contained in this Fact Sheet.

Whilst CANSA has taken every precaution in compiling this Fact Sheet, neither it, nor any contributor(s) to this Fact Sheet can be held responsible for any action (or the lack thereof) taken by any person or organisation wherever they shall be based, as a result, direct or otherwise, of information contained in, or accessed through, this Fact Sheet.



Sources and References Consulted or Utilised

Breast Cancer Fund

<http://www.breastcancerfund.org/clear-science/radiation-chemicals-and-breast-cancer/ionizing-radiation.html?referrer=https://www.google.co.za/>

Brenner, D. J. 2011. Are x-ray backscatter scanners safe for airport passenger screening? For most individuals, probably yes, but a billion scans per year raises long-term public health concerns. *Radiology*, 259(1), 6–10.

Electromagnetic Spectrum

https://altered-states.net/cart/p2070/3-AXIS_EMF_RF_Radiation_ElectroSmog_Meter

Goodman, T.R.

<http://www.imagewisely.org/imaging-modalities/computed-tomography/imaging-physicians/articles/ionizing-radiation-effects-and-their-risk-to-humans>

Image Wisely

<http://www.imagewisely.org/imaging-modalities/computed-tomography/imaging-physicians/articles/ionizing-radiation-effects-and-their-risk-to-humans>

Medscape

<http://www.medscape.com/viewarticle/824999>

Mirion Technologies

<https://www.mirion.com/introduction-to-radiation-safety/types-of-ionizing-radiation/>

RadiologyInfo

<http://www.radiologyinfo.org/en/info.cfm?pg=safety-xray>

Understanding Medical Radiation

<https://www.medicalradiation.com/facts-about-radiation/benefits-and-risks-of-radiation/benefits-and-risks-of-ionizing-radiation/>

Wikipedia

<https://en.wikipedia.org/wiki/Becquerel>

World Health Organization

<http://www.who.int/mediacentre/factsheets/fs371/en/>

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Diagnostic Radiographer; Dip Audiometry and Noise Measurement; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

October 2018