

Cancer Association of South Africa (CANSA)



Research • Educate • Support

Fact Sheet on Clear-Cell Adenosarcoma of the Vagina

Introduction

Clear-cell adenocarcinoma (CCA) of the vagina (or cervix) is a rare cancer often linked to the drug diethylstilbestrol (DES), which was prescribed in the mistaken belief that it prevented miscarriage and ensured a healthy pregnancy.

[Picture Credit: Diethylstilbestrol]




This synthetic oestrogen was given to millions of pregnant women, primarily from 1938-1971 but not limited to those years. Internationally, DES use continued until the early 1980s. DES was given if a woman had a previous miscarriage, diabetes, or a problem-pregnancy with bleeding, threatened miscarriage or premature labour. Up until the mid to late 1950s some women were given DES shots. After that, DES was primarily prescribed in pill form. DES also was included in some prenatal vitamins. In the late 1960s through 1971 a cluster of young women, from their teens into their twenties, was mysteriously diagnosed with CCA, a cancer not generally found in women until after menopause.

Clear-Cell Adenosarcoma (CCA) of the Vagina

Clear-Cell Adenosarcoma (CCA) of the vagina is one of the most common subtype of vaginal adenocarcinoma associated with diethylstilbestrol (DES) exposure in young females. CCA of the vagina can also occur in postmenopausal women without exposure to DES. It is a rare vaginal cancer, accounting for 5% to 10% of primary vaginal malignancies (PathologyOutlines.com).

[Picture Credit: Female Anatomy]



© 2008 Ramin Mirhashemi, MD, www.gynla.com

The relative risk of CCA of the Vagina in DES Daughters

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Dip Audiometry and Noise Measurement; Diagnostic Radiographer; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

September 2018

(the daughter of a woman who received diethylstilbestrol (DES) during pregnancy) is much higher compared to the general population.

Incidence of Clear-Cell Adenosarcoma of the Vagina

The National Cancer Registry (2014) does not provide any information regarding Clear-cell Adenosarcoma of the Vagina. According to the National Cancer Registry (2014) the following number of cancer of the vagina cases was histologically diagnosed in South Africa during 2014:

Group - Females 2014	Actual No of Cases	Estimated Lifetime Risk	Percentage of All Cancers
All females	164	1:1 520	0,43%
Asian females	5	1:1 876	0,43%
Black females	127	1:1 481	0,79%
Coloured females	12	1:1 705	0,30%
White females	20	1:1 774	0,12%

The frequency of histologically diagnosed cases of cancer of the vagina in South Africa for 2014 was as follows (National Cancer Registry, 2014):

Group - Females 2014	0 – 19 Years	20 – 29 Years	30 – 39 Years	40 – 49 Years	50 – 59 Years	60 – 69 Years	70 – 79 Years	80+ Years
All females	1	5	22	35	38	32	14	8
Asian females	0	0	0	1	0	1	1	0
Black females	1	4	21	32	26	20	10	7
Coloured females	0	1	1	0	6	2	0	0
White females	0	0	0	2	6	8	3	1

N.B. In the event that the totals in any of the above tables do not tally, this may be the result of uncertainties as to the age, race or sex of the individual. The totals for 'all males' and 'all females', however, always reflect the correct totals.

Risk Factors for Clear-Cell Adenosarcoma of the Vagina

Age and being exposed to the drug DES (diethylstilbestrol) before birth affect a woman's risk of vaginal cancer.

Anything that increases one's risk of getting a disease is called a risk factor. Having a risk factor does not mean that one will get cancer; not having risk factors does not mean that one will not get cancer. Talk with a doctor if you think you may be at risk. Risk factors for vaginal cancer include the following:

- Being aged 60 or older.
- Being exposed to DES while in the mother's womb. In the 1950s, the drug DES was given to some pregnant women to prevent miscarriage (premature birth of a foetus that cannot survive). Women who were exposed to DES before birth have an increased risk of vaginal cancer. Some of these women develop a rare form of vaginal cancer called clear cell adenocarcinoma.
- Having human papilloma virus (HPV) infection.
- Having a history of abnormal cells in the cervix or cervical cancer.
- Having a history of abnormal cells in the uterus or cancer of the uterus.
- Having had a hysterectomy for health problems that affect the uterus.

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Dip Audiometry and Noise Measurement; Diagnostic Radiographer; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

September 2018

Signs and Symptoms of Clear-Cell Adenosarcoma of the Vagina

Patients may have one or more of the following:

Vaginal bleeding

Dyspareunia (painful sexual intercourse)

(Hanselaar, *et al.*)

Diagnosis of Clear-cell Adnosarcoma of the Vagina

The following tests and procedures may be used:

Physical examination and history: Physical examination including a history of the patient's health habits and past illnesses and treatments.

Pelvic examination: an examination of the vagina, cervix, uterus, fallopian tubes, ovaries, and rectum.

Pap smear: A procedure to collect cells from the surface of the cervix and vagina.

Biopsy: The removal of cells or tissues from the vagina and cervix so they can be viewed under a microscope by a pathologist to check for signs of cancer. A biopsy may be done during a colposcopy.

Colposcopy: A procedure in which a colposcope (a lighted, magnifying instrument) is used to check the vagina and cervix for abnormal areas. Tissue samples may be taken using a curette (spoon-shaped instrument) and checked under a microscope for signs of disease.

Treatment of Clear-Cell Adenosarcoma of the Vagina

Approximately 5% of primary vaginal malignancies are adenocarcinomas. Whenever this diagnosis is considered, it is necessary to rule out metastatic lesions from the bowel, uterus, or ovary. The most common variant is the clear-cell adenocarcinoma, which can occur spontaneously and in women with *in utero* exposure to DES. Primary non-DES-related adenocarcinoma of the vagina is rare and occurs predominately in postmenopausal women.

Treatment may involve surgical intervention and radiation therapy. For stage I clear-cell adenocarcinoma in the typical young patient, surgery may be considered to preserve ovarian function. Surgery for vaginal clear-cell adenocarcinoma usually requires a radical hysterectomy and vaginectomy with reconstruction. The vaginectomy is performed only to the level required to obtain an adequate margin. Local excision appears inferior to radical surgery. The role of chemotherapy has not been determined.

About Clinical Trials

Clinical trials are research studies that involve people. They are conducted under controlled conditions. Only about 10% of all drugs started in human clinical trials become an approved drug.

Clinical trials include:

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Dip Audiometry and Noise Measurement; Diagnostic Radiographer; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

September 2018

- Trials to test effectiveness of new treatments
- Trials to test new ways of using current treatments
- Tests new interventions that may lower the risk of developing certain types of cancers
- Tests to find new ways of screening for cancer

The [South African National Clinical Trials Register](#) provides the public with updated information on clinical trials on human participants being conducted in South Africa. The Register provides information on the purpose of the clinical trial; who can participate, where the trial is located, and contact details.

For additional information, please visit: www.sanctr.gov.za/

Medical Disclaimer

This Fact Sheet is intended to provide general information only and, as such, should not be considered as a substitute for advice, medically or otherwise, covering any specific situation. Users should seek appropriate advice before taking or refraining from taking any action in reliance on any information contained in this Fact Sheet. So far as permissible by law, the Cancer Association of South Africa (CANSA) does not accept any liability to any person (or his/her dependants/**estate/heirs**) relating to the use of any information contained in this Fact Sheet.

Whilst the Cancer Association of South Africa (CANSA) has taken every precaution in compiling this Fact Sheet, neither it, nor any contributor(s) to this Fact Sheet can be held responsible for any action (or the lack thereof) taken by any person or organisation wherever they shall be based, as a result, direct or otherwise, of information contained in, or accessed through, this Fact Sheet.



Sources and References Consulted or Utilised

Center for Disease Control and Prevention

http://www.cdc.gov/des/hcp/information/daughters/risks_daughters.html

Hanselaar, A., van Loosbroek, M., Schuurbijs, O., Helmerhorst, T., Bulten, J. & Bernheim, J. 2000. Clear cell adenocarcinoma of the vagina and cervix.

[https://doi.org/10.1002/\(SICI\)1097-0142\(19970601\)79:11<2229::AID-CNCR22>3.0.CO;2-X](https://doi.org/10.1002/(SICI)1097-0142(19970601)79:11<2229::AID-CNCR22>3.0.CO;2-X)

National Cancer Institute

<http://www.cancer.gov/cancertopics/pdq/treatment/vaginal/Patient/page1>

<http://www.cancer.gov/clinicaltrials/learningabout/what-are-clinical-trials>

<http://www.cancer.gov/about-cancer/treatment/clinical-trials>

NCBI

<http://www.ncbi.nlm.nih.gov/books/NBK13130/>

Oncolink

<http://www.oncolink.org/types/article.cfm?c=215&id=9501>

PathologyOutlines.com

<http://www.pathologyoutlines.com/topic/vaginalclearcelladeno.html>

Researched and Authored by Prof Michael C Herbst

[D Litt et Phil (Health Studies); D N Ed; M Art et Scien; B A Cur; Dip Occupational Health; Dip Genetic Counselling; Dip Audiometry and Noise Measurement; Diagnostic Radiographer; Medical Ethicist]

Approved by Ms Elize Joubert, Chief Executive Officer [BA Social Work (cum laude); MA Social Work]

September 2018

Patient.co.uk

<http://www.patient.co.uk/doctor/vaginal-cancer>

The Global Library of Women's Medicine

http://www.glowm.com/section_view/heading/Vaginal%20Cancer/item/260

WebMD

<http://www.webmd.com/cancer/tc/vaginal-cancer-treatment-patient-information-nci-pdq-general-information-about-vaginal-cancer>

Wikipedia

http://en.wikipedia.org/wiki/Clear-cell_adenocarcinoma_of_the_vagina